

**Galveston Bay Status and Trends Project  
2004-2006**

**Final Report**



**Galveston Bay  
Status and Trends**

**Contract Number 582-4-65047**

**2006**

**Prepared For:**

Texas Commission on Environmental Quality  
Galveston Bay Estuary Program  
17041 El Camino Real, Ste. 210  
Houston, Texas 77058

Project Manager  
Steven R. Johnston  
[SJohnsto@tceq.state.tx.us](mailto:SJohnsto@tceq.state.tx.us)



**Prepared By:**

Geotechnology Research Institute (GTRI)  
Houston Advanced Research Center (HARC)  
4800 Research Forest Drive  
The Woodlands, Texas 77381

Principal Investigator  
L. James Lester  
[jlester@harc.edu](mailto:jlester@harc.edu)

Project Manager  
Lisa A. Gonzalez  
[lgonzalez@harc.edu](mailto:lgonzalez@harc.edu)



*Prepared in Cooperation with the  
Texas Commission on Environmental Quality and  
U.S. Environmental Protection Agency*

*The preparation of this report was  
financed through grants from the  
U.S. Environmental Protection Agency through the  
Texas Commission on Environmental Quality*

[www.galvbaydata.org](http://www.galvbaydata.org)

# Table of Contents

<b>1</b>	<b>Executive Summary</b>	<b>9</b>
<b>2</b>	<b>Introduction</b>	<b>15</b>
<b>3</b>	<b>Methodology and Results</b>	<b>18</b>
<b>3.1</b>	<b>Water and Sediment Quality</b>	<b>18</b>
3.1.1	Recent Trends in Water and Sediment Quality (1999-2005)	20
3.1.2	Water Quality Gradients in Clear Creek and Clear Lake	32
<b>3.2</b>	<b>Coastal Fisheries</b>	<b>46</b>
3.2.1	Length of Commercial and Recreational Species	46
3.2.2	Biodiversity Analysis of Coastal Fisheries Resources	58
<b>3.3</b>	<b>Habitat</b>	<b>77</b>
3.3.1	Assessment of Wetland Loss and Gain Projections	77
3.3.2	Habitat Fragmentation Analysis	92
<b>3.4</b>	<b>Data Gaps Assessment</b>	<b>101</b>
3.4.1	Water and Sediment Quality	101
3.4.2	Living Resources	106
3.4.3	Habitat	108
<b>3.5</b>	<b>Other Indicators of Bay Health</b>	<b>110</b>
3.5.1	Seafood Safety	110
3.5.2	Colonial Nesting Waterbirds	116
3.5.3	Galveston Bay Shoreline Analysis	118
<b>3.6</b>	<b>Outreach</b>	<b>121</b>
3.6.1	Charting the Course Issue Papers	121
3.6.2	Website Development: <a href="http://www.galvbaydata.org">www.galvbaydata.org</a>	121
<b>4</b>	<b>Conclusion and Lessons Learned</b>	<b>125</b>
<b>5</b>	<b>Bibliography</b>	<b>126</b>
<b>6</b>	<b>Appendices</b>	<b>127</b>

# Tables and Figures

<b>Tables</b>	<b>Page</b>
<b>Water and Sediment Quality</b>	
Table 3.1.1. TCEQ screening levels for nutrients in surface waters of tributaries (tidal stream screening levels) and subbays (estuarine screening levels).	20
Table 3.1.2. Indicator rating system developed for the nutrients in surface waters indicator.	20
Table 3.1.3. Indicator describing the state of nutrients (ammonia, nitrate-nitrite, total phosphorus) and chlorophyll-a concentrations in surface waters of the Lower Galveston Bay watershed.	21
Table 3.1.4. TCEQ screening levels for nutrients in surface waters of tributaries (tidal stream screening levels) and subbays (estuarine screening levels).	25
Table 3.1.5. Indicator rating system developed for the bacterial pathogens in surface waters. Table created by the Galveston Bay Indicators Project (2005) based on screening levels from the TCEQ (2004).	25
Table 3.1.6. Indicator describing the state of bacterial pathogens in surface waters as a proportion of the TCEQ screening level.	26
Table 3.1.7. Indicator rating system developed for toxic contaminants in sediments.	29
Table 3.1.8. Organic pesticides and industrial organics in sediment as a proportion of Marine Probable Effects Levels (PELs) published by the TCEQ.	29
Table 3.1.9. Detail of industrial organics in sediments of the Houston Ship Channel.	30
Table 3.1.10. Heavy metals in sediment as a proportion of Marine Probable Effects Levels (PELs) published by the TCEQ (2004).	30
Table 3.1.11. Detail of nine heavy metals in sediments of the Houston Ship Channel.	31
Table 3.1.12. Average chlorophyll-a and pheophytin-a concentrations exceeding the surface water quality screening level of 19.2 ug/L at ten stations in the Clear Creek watershed.	41
Table 3.1.13. Twenty-six organic and inorganic sediment quality parameters analyzed for stations in Clear Creek/Clear Lake.	45
<b>Coastal Fisheries</b>	
Table 3.2.1. Correlations between environmental variables associated with bag seine collections.	47
Table 3.2.2. List of species selected for analysis from bag seine data, the number included in the sample and the average length in mm of the sampled individuals.	47
Table 3.2.3. Correlation between lengths of selected species collected in bag seine and water quality variables.	48
Table 3.2.4. Results of ANOVA analysis of temporal variation in log transformed length data of six selected species from bag seine collections.	48
Table 3.2.5. Correlations between environmental variables associated with shrimp trawl collections.	49
Table 3.2.6. List of species selected for analysis from shrimp trawl data, the number included in the sample and the average length in mm of the sampled individuals.	49
Table 3.2.7. Correlation between length of selected species collected in shrimp trawls and water quality variables	50

## Tables Continued...

	<b>Page</b>
Table 3.2.8. Results of ANOVA analysis of temporal variation in log transformed length data of six selected species from shrimp trawl collections.	51
Table 3.2.9. Species of commercially and recreationally imported fish analyzed for two gear types.	52
Table 3.2.10. Indicator describing trends in length of commercially and recreationally import species of fish captured in gill net and shrimp trawl.	57
Table 3.2.11. Number of sampling events for each gear type in each subbay of the Galveston Bay System used in calculating biodiversity.	59
Table 3.2.12. Lists of the common names of the 25 species captured by bag seine in the five subbays that have the highest ranks in relative abundance over the period of record from 1977 to 2004.	62
Table 3.2.13. Spearman rank order correlations of the ranked relative abundances of species captured by bag seine in the five subbays of Galveston Bay.	63
Table 3.2.14. Lists of the common names of the 25 species captured by shrimp trawl in four subbays that have the highest ranks in relative abundance over the period of record from 1982 to 2004.	64
Table 3.2.15. Spearman Rank Order Correlations of the Ranked Relative Abundances of Species Captured by Shrimp Trawl in the Four Subbays.	65
Table 3.2.16. Lists of the common names of the 25 species captured by gill net in five subbays that have the highest ranks in relative abundance over the period of record from 1977 to 2004.	66
Table 3.2.17. Spearman rank order correlations between the species captured in gill net samples from the five subbays ranked by average relative abundance over the period of record.	67
Table 3.2.18. Monthly mean S-W Index values over the period of record from 1977 to 2004 based on bag seine collections from five subbays.	67
Table 3.2.19. Monthly mean S-W Index values based on shrimp trawl samples collected in four subbays over the period of record from 1982 to 2004.	68
Table 3.2.20. Monthly mean S-W Index values based on gill net samples collected in five subbays over the period of record from 1977 to 2004.	69
Table 3.2.21. Annual average S-W Index values for bag seine samples from five subbays over the period of 1978 to 2004.	70
Table 3.2.22. Bag Seine Biodiversity by Subbay based on annualized samples.	71
Table 3.2.23. Annual Shannon-Wiener Index for Shrimp Trawl Samples from Four Subbays for the Period of Record; 1982-2004.	72
Table 3.2.24. Gill net biodiversity by year in all subbays of Galveston Bay based on average S-W values from individual collections.	73
Table 3.2.25. S-W Biodiversity values calculated using annualized relative abundances for gill net collections.	74
Table 3.2.26. Average over all years for all subbays based on the annualized S-W Index for all gear types.	75

## Habitat

Table 3.3.1. Change in acreage of five wetland classes in 21 subwatersheds of the Lower Galveston Bay watershed, 1996 to 2001.	81
Table 3.3.2. Change in acreage of wetlands in 21 watersheds of the Lower Galveston Bay watershed, 1996 to 2001.	82

## Tables Continued...

	<b>Page</b>
Table 3.3.3. Change in acreage of five wetland classes in five counties of the Lower Galveston Bay watershed, 1996 to 2001.	83
Table 3.3.4. Change in acreage of wetlands in 5 counties of the Lower Galveston Bay watershed, 1996 to 2001.	83
Table 3.3.5. Projected change in acreage of five wetland classes in 21 subwatersheds of the Lower Galveston Bay watershed, 1996 to 2016.	84
Table 3.3.6. Projected change in acreage of wetlands in 21 watersheds of the Lower Galveston Bay watershed, 1996 to 2016.	85
Table 3.3.7. Projected change in acreage of five wetland classes in five counties of the Lower Galveston Bay watershed, 1996 to 2016.	86
Table 3.3.8. Projected change in acreage of wetlands in 5 counties of the Lower Galveston Bay watershed, 1996 to 2016.	86
Table 3.3.9. List of Galveston Bay Sub Watersheds with the Fragmentation Indices Calculated for Wetland and Woody Land Cover Classifications.	97

## Data Gaps Assessment

No tables in this section -

## Other Indicators of Bay Health

Table 3.5.1. Rating of risk from consumption of contaminants in fish captured in various parts of the Galveston Bay system.	113
Table 3.5.2. Rating of water quality with regards to fecal coliform bacteria in the major subbays and tributaries of Galveston Bay over the last four decades.	114
Table 3.5.3. Indicator describing the twenty-year trend in abundance of colonial waterbird nesting pairs.	117

## Figures

## Page

### Water and Sediment Quality

Figure 3.1.1. Tributaries and subbays of Galveston Bay per the Galveston Bay Segmentation Scheme as modified by the Status and Trends Project in 2003.	19
Figure 3.1.2. Stations sampled for nutrients and chlorophyll-a in surface waters of Galveston Bay 1999-2005. Stations are color-coded according to the proportion of samples exceeding TCEQ screening levels.	22
Figure 3.1.3. Monthly Average Dissolved Oxygen Concentrations in Surface Waters of Galveston Bay and Tributaries; 1969-2003.	23
Figure 3.1.4. Stations where dissolved oxygen was collected from 5:00 am to 10:00 am in surface waters of Galveston Bay 1999-2005.	24
Figure 3.1.5. Stations sampled for pathogens (fecal coliform bacteria, E. coli, and Enterococci) in surface waters of Galveston Bay 1999-2005.	27
Figure 3.1.6. Description of the Area, Species, and Contaminants Included in Galveston Bay Seafood Advisories Issued by the Texas Department of State Health Services in 1990, 2201, and 2005.	31

## Figures Continued...

	<b>Page</b>
Figure 3.1.7. TCEQ Surface Water Quality Monitoring Program (SWQM) and Clean Rivers Program (CRP) monitoring stations identified in tidally influenced waters of Clear Creek watershed.	33
Figure 3.1.8. Average salinities by station ( $n \geq 12$ ) in surface waters of Clear Creek and Clear Lake as sampled by the TCEQ from 1999-2004.	34
Figure 3.1.9. Average BOD5 by station ( $n \geq 12$ ) in surface waters of Clear Creek and Clear Lake as sampled by the TCEQ from 1999-2004.	35
Figure 3.1.10. Average dissolved oxygen (DO) by station ( $n \geq 12$ ) in surface waters of Clear Creek and Clear Lake as sampled by the TCEQ from 1999-2004.	36
Figure 3.1.11. Average concentrations of nitrate-nitrite by station ( $n \geq 12$ ) in surface waters of Clear Creek and Clear Lake as sampled by the TCEQ from 1999-2004.	37
Figure 3.1.12. Average concentrations of total phosphorus, dissolved phosphorus, and orthophosphate by station ( $n \geq 12$ ) in surface waters of Clear Creek and Clear Lake as sampled by the TCEQ from 1999-2004.	38
Figure 3.1.13. Average concentrations of chlorophyll-a and pheophytin-a by station ( $n \geq 12$ ) in surface waters of Clear Creek and Clear Lake as sampled by the TCEQ from 1999-2004.	40
Figure 3.1.14. Average concentrations of fecal coliform bacteria by station ( $n \geq 12$ ) in surface waters of Clear Creek and Clear Lake as sampled by the TCEQ from 1999-2004.	42
Figure 3.1.15. Average concentrations of <i>E. coli</i> by station ( $n \geq 12$ ) in surface waters of Clear Creek and Clear Lake as sampled by the TCEQ from 1999-2004.	43
Figure 3.1.16. Average concentrations of Enterococci by station ( $n \geq 12$ ) in surface waters of Clear Creek and Clear Lake as sampled by the TCEQ from 1999-2004.	44

## Coastal Fisheries

Figure 3.2.1. Trends in annual average length of (a) Red drum in gill net, (b) sheepshead in gill net, (c) Southern flounder in gill net, and (d) spotted seatrout in gill net.	53
Figure 3.2.2. Trends in annual average length of (a) Atlantic croaker in shrimp trawl, (b) spotted seatrout in shrimp trawl, and (c) white shrimp in shrimp trawl.	55
Figure 3.2.3. Trends in Annual Average S-W Index values over the period of record from 1977 to 2004 based on collections from Upper and Lower Galveston Bay.	76

## Habitat

Figure 3.3.1. Map depicting geographic differences between the counties and subwatersheds existing in the Lower Galveston Bay watershed.	80
Figure 3.3.2. Number of approved COE Section 10/404 wetland permits issued in the five counties surrounding Galveston Bay during the period 1992-2003.	88
Figure 3.3.3. Number of wetland permits issued by the US Army Corps of Engineers Galveston District in the five counties surrounding Galveston Bay during the period 1992-2003.	89
Figure 3.3.4. Number of wetland permits issued by the US Army Corps of Engineers Galveston District in 21 subwatersheds surrounding Galveston Bay during the period 1992-2003.	90
Figure 3.3.5. Number of wetland permits issued by the US Army Corps of Engineers Galveston District by permit type during the period 1992-2003.	91

## Figures Continued...

	<b>Page</b>
Figure 3.3.6. Classified image of the five counties using the modified classification scheme.	93
Figure 3.3.7. County-level wetland fragmentation of the Lower Galveston Bay watershed.	95
Figure 3.3.8. County-level woody land fragmentation of the Lower Galveston Bay watershed.	96
Figure 3.3.9. Subwatershed-level wetland land fragmentation of the Lower Galveston Bay watershed based on the habitat index ( <i>F</i> ).	98
Figure 3.3.10. Subwatershed-level woody land fragmentation of the Lower Galveston Bay watershed based on the habitat index ( <i>F</i> ).	99

## Data Gaps Assessment

Figure 3.4.1. Total number of samples by area of three classes of toxic contaminants sampled in sediments of Galveston Bay during the period 1973-2004	102
Figure 3.4.2. Total number of samples by year of three classes of toxic contaminants sampled in sediments of Galveston Bay during the period 1973-2004.	103
Figure 3.4.3. Total number of samples by area of three classes of water quality parameters sampled in surface waters of Galveston Bay during the period 1969-2004.	104
Figure 3.4.4. Total number of samples by year of three classes of water quality parameters sampled in surface waters of Galveston Bay during the period 1969-2004.	105
Figure 3.4.5. Total number of TPWD gill net samples by month in Galveston Bay over the period 1976-2004.	107

## Other Indicators of Bay Health

Figure 3.5.1. Comparison of DSHS Approved, Conditionally Approved, and Restricted Shellfish Harvest Areas in Galveston Bay for the Years 1996, 2000, and 2003.	115
Figure 3.5.2. Increasing twenty year trend in abundance of brown pelican nesting pairs in Galveston Bay.	118
Figure 3.5.3. Developed and undeveloped shoreline of Galveston Bay.	120

## Website Development: [www.galvbaydata.org](http://www.galvbaydata.org)

Figure 3.6.1. Water and Sediment Quality Data Portal query result as seen at <a href="http://www.galvbaydata.org">www.galvbaydata.org</a> .	122
Figure 3.6.2. Fisheries Data Portal query result as seen at <a href="http://www.galvbaydata.org">www.galvbaydata.org</a> .	124